

Hadronic few-body systems in chiral dynamics

Few-body systems in hadron physics

Daisuke Jido

Received: date / Accepted: date

Abstract Hadronic composite states are introduced as few-body systems in hadron physics. The $\Lambda(1405)$ resonance is a good example of the hadronic few-body systems. It has turned out that $\Lambda(1405)$ can be described by hadronic dynamics in a modern technology which incorporates coupled channel unitarity framework and chiral dynamics. The idea of the hadronic $\bar{K}N$ composite state of $\Lambda(1405)$ is extended to kaonic few-body states. It is concluded that, due to the fact that K and N have similar interaction nature in s -wave \bar{K} couplings, there are few-body quasibound states with kaons systematically just below the break-up thresholds, like $\bar{K}NN$, $\bar{K}KN$ and $\bar{K}KK$, as well as $\Lambda(1405)$ as a $\bar{K}N$ quasibound state and $f_0(980)$ and $a_0(980)$ as $\bar{K}K$.

Keywords Hadronic composite state · $\Lambda(1405)$ resonance · kaonic few-body systems · chiral dynamics

1 Introduction

Hadrons are composite objects of quark and gluons governed by quantum chromodynamics, QCD. So far hundreds of hadrons have been experimentally observed in spite of only the five (or six) fundamental pieces, up, down, strange, charm, bottom (and top) quarks. The richness of hadron spectrum is a consequence of highly nontrivial dynamics of quarks and gluons confined in hadron. Precise measurement of hadron spectrum is one of the phenomenological ways to view colored dynamics of quarks and gluons inside hadron, and gives us clues of the confinement mechanism. Especially the systematics of energy excitation is a key issue to understand the hadron structure.

A conventional picture of hadron is the quark model, in which one considers constituent quarks as quasi-particles moving in a one-body mean field created

D. Jido
Yukawa Institute for Theoretical Physics Kyoto University, Kyoto 606-8502, JAPAN
Tel.: +81-75-7537034
Fax: +81-75-7537010
E-mail: jido@yukawa.kyoto-u.ac.jp

by gluon dynamics, and symmetry of the quasi-particles and (weak) residual interaction characterize the hadron structure. Although quark models work well for lowest-lying states, there are a lot of states found which do not match quark model predictions, which are so-called exotic hadrons, and quark models should incorporate hadron dynamics once strong decay channels open. It has been realized that quark models do not give us the universal picture of the hadron structure, and that the constituent quark alone cannot be an effective degree of freedom in hadron. These facts bring us the idea to consider other effective constituents, such as diquark correlation, multi-quark component and hadronic composite, which will be the origin of richness of hadron spectrum.

2 Hadronic composite states

Hadron composite state can be one of the existence forms of hadron. Its constituents are hadrons governed by hadronic dynamics not inter-quark colored force. Consequently the spacial size is much larger than the typical hadron size which is characterized by the confining force. The typical examples are atomic nuclei which are bound states of nucleons. Mesons also can be constituents of the hadron composite states. Because of no number conservation of mesons, the composite state with mesons has transition modes to lighter mesons, mostly pions, and absorptive decay modes. Therefore, most of hadronic composite states are unstable bound states with decay modes.

Hadron composite state can be a new sample to investigate the confinement mechanism. Most of hadron flavors can be explained by a quark-antiquark pair $\bar{q}q$ and three quarks qqq . So far, very few flavor exotic hadrons have been observed. This is an important question to be solved in hadron physics. A clue is multi-quark state. Multi-quark state always has color singlet clusters. Thus, there is competition between colorful hadron *constituent force* inside confinement range and colorless hadron *interaction force* in a larger range than the typical hadron size. For instance, the H -dibaryon composed by $uuddss$ can have a colored three diquark configuration ($[ud][ds][su]$) governed by short range color interaction and a colorless two baryon configuration ($[AA]$ and/or $[\Xi N]$) governed by longer range hadronic interaction. The real state can be a mixture of these configurations. Nevertheless, since the former configuration should be a compact object, while the latter can be a larger object, there is a (small) scale gap between two configurations. These two have also different systematics in excitation modes and flavor symmetry. Thus, it would be very interesting if one could know which systematics is realized in each hadron.

3 The $\Lambda(1405)$ resonance

$\Lambda(1405)$ is a baryon resonance with isospin $I = 0$ and strangeness $S = -1$ sitting between the $\bar{K}N$ (1435 MeV) and $\pi\Sigma$ (1331 MeV) thresholds with mass around 1405 MeV and 50 MeV decay width to $\pi\Sigma$. The flavor of $\Lambda(1405)$ can be expressed by the minimal quark contents of uds , however, simple quark models have failed to reproduce the $\Lambda(1405)$ mass. $\Lambda(1405)$ is a historical example of the hadronic composite state. $\Lambda(1405)$ has been considered as a quasibound state of $\bar{K}N$ [1, 2], even before QCD was established. In general, the physically observed state should

be given as a mixture of the hadronic composite and quark model type components, it is important to examine the dominant components of hadron resonance states to understand their structure and dynamics, and it would be interesting if one could know the fraction of the components.

For the theoretical description of $\Lambda(1405)$, one needs dynamical study of coupled channels including at least $\bar{K}N$ and $\pi\Sigma$, because the $\Lambda(1405)$ resonance is located in the 100 MeV window of the $\bar{K}N$ and $\pi\Sigma$ thresholds. Flavor SU(3) complete treatment of coupled channels, by including $\pi\Sigma$, $\bar{K}N$, $\pi\Lambda$, $\eta\Lambda$, $\eta\Sigma$ and $K\Sigma$, was already done using phenomenological vector-meson exchange potential in Ref. [3]. Coupled channels calculation of $\pi\Sigma$ and $\bar{K}N$ based on the cloudy bag model was done in Ref. [4]¹. The modern approach based on chiral dynamics and unitary coupled channels was initiated by Ref. [5], and the $\Lambda(1405)$ resonance is well described as well as the K^-p threshold properties and K^-p scatterings. (See Ref. [6] as a recent review article and Ref. [7, 8] for a very recent update including the new SIDDHARTA measurement [9].)

One of the most important consequences of the coupled channels approach based on chiral dynamics is that the $\Lambda(1405)$ resonance is a superposition of two states [10, 11]. These two states have different properties [11]: one of the states is located at 1426 MeV with a 32 MeV width and couples dominantly to $\bar{K}N$, while the other is at 1390 MeV with a broader width and couples strongly to $\pi\Sigma$. All of the recent calculations based on chiral interaction suggest the two pole structure of $\Lambda(1405)$ although the pole positions are model dependent slightly. The reason that there are two states around the $\Lambda(1405)$ energy region is that the chiral interaction indicates two attractions with $I = 0$ in the flavor SU(3) singlet and octet channels group-theoretically [11] or in the $\bar{K}N$ and $\pi\Sigma$ channels physically [12]. The latter fact implies that the $\Lambda(1405)$ is essentially described by two dynamical channels of $\bar{K}N$ and $\pi\Sigma$, and that the higher state is a $\bar{K}N$ quasibound state decaying to $\pi\Sigma$ and the lower is a $\pi\Sigma$ resonant state [12]. As a consequence of the double pole nature, the $\Lambda(1405)$ spectrum depends on the initial channel [11]. Thus, observing that the $\Lambda(1405)$ spectrum in the $\bar{K}N \rightarrow \pi\Sigma$ channel is different from that in $\pi\Sigma \rightarrow \pi\Sigma$, one can confirm that $\Lambda(1405)$ is a dynamical object of $\bar{K}N$ and $\pi\Sigma$. For this purpose, since $\Lambda(1405)$ is located below the threshold of $\bar{K}N$, one needs indirect production of $\Lambda(1405)$ from $\bar{K}N$. It has been found in Refs. [13, 14] that in the $K^-d \rightarrow \Lambda(1405)n$ reaction $\Lambda(1405)$ is selectively produced by $\bar{K}N$, thus this reaction is one of the suitable reactions to investigate the nature of $\Lambda(1405)$. An experiment of this reaction at J-PARC is proposed [15] observing neutrons in the forward direction.

4 Compositeness of hadron

Compositeness of hadron had been discussed before QCD was established in order to find fundamental or elementary hadrons. But this trial was certainly failed, since all of the hadrons are composite objects of quarks and gluons. Here we discuss the compositeness of hadron resonances in terms of quark originated state and hadronic composite state. To discuss the compositeness of the hadron resonances, first of all, we need to define elementary hadrons. The elementary hadrons

¹ In this model, a bare pole term for the $\Lambda(1405)$ resonance was explicitly introduced around 1650 MeV.

may be the ground state hadrons stable against strong interactions, or one may take hadrons which survive in the large N_c limit. Once one defines the elementary hadrons, one can discuss the compositeness of hadron quantitatively. The elementary components are expressed in the free Hamiltonian H_0 and the composite states are dynamically generated as a consequence of the interaction of the elementary components V out of the full Hamiltonian of the system $H = H_0 + V$. The compositeness index can be written as $1 - Z$ where $Z = \sum_n |\langle n|d \rangle|^2$ with the eigenstates $|n\rangle$ of H_0 and the dynamically generated state $|d\rangle$ as an eigenstate of H [16]. This definition of the compositeness can be extended in field theory by introduction the Lagrangian of the system as $L = L_0 + L_{\text{int}}$ in which L_0 is the free Lagrangian of the elementary component and L_{int} represents their interaction. The field renormalization constant Z is shown in a residue of the full propagator Δ satisfying Dyson equation at the pole position, of which normalization is given by the free propagator $\Delta_0(E) = 1/(E - M_0)$ with the bare mass M_0 [17]. An application for chiral dynamics is discussed in Ref. [18]. For the resonance state, the compositeness index can be a complex number. Thus, one needs its appropriate interpretation, which is an open issue yet.

Here we would like to discuss the compositeness of $\Lambda(1405)$ in a different way within the chiral unitary approach [19]. In the chiral unitary approach, one solves Lippmann-Schwinger equation $T = V + VGT$, in which V is the interaction kernel, while G is the loop function which guarantees unitarity and specifies the model space. Thus, in the chiral unitary approach, the elementary components are given in the loop function. In the present case for $\Lambda(1405)$, they are the lowest lying octet baryons and mesons. The interactions among the elementary components are given by the chiral effective theory. In the context of the discussion of compositeness, one has to take care of the interaction kernel, since in the interaction kernel sources of the resonances can be hidden. It is well-known that the s -channel resonance contributions are involved in the contact interactions of the elementary component [20]. Nevertheless, the Weinberg-Tomozawa interaction may be free from the s -wave resonance contributions since it comes from the t -channel vector meson exchange. For the discussion of the compositeness of the resonance, let us take only the Weinberg-Tomozawa interaction as the interaction kernel. Since the Weinberg-Tomozawa interaction is a short range contact force, one needs to regularize the loop integral. In the regularization procedure, one fixes high-momentum behavior which is not controlled in the present model space. Therefore, even though one takes only hadronic interaction in interaction kernel, some contributions coming from outside of the model space can be hidden in the regularization parameters [19]. In the chiral unitary model, this parameter is fixed phenomenologically so as to reproduce observed scattering cross section. Thus, if necessary, nature will request the components which come from the outside of the model space, namely, quark originated component.

We have a choice of the parameter in which the hidden contribution can be excluded from formulation by theoretical requirement on the renormalization constant (natural renormalization scheme) [19]. If resonances can be reproduced by the chiral unitary approach with the Weinberg-Tomozawa interaction in the natural renormalization scheme, the resonances can be regarded as composite objects of meson and baryon constituents. In Ref. [19] it has been found that the pole positions of $\Lambda(1405)$ is well reproduced by the natural renormalization scheme, while $N(1535)$ is not. This suggests that $\Lambda(1405)$ is mostly a composite state of




Fig. 1 Transition form factors of $\gamma^* p \rightarrow N(1535)$ calculated by the chiral unitary model (meson cloud) [22] and spectator quark model (valence quark) [23,24]. The plots are taken from Ref. [21].

the ground state mesons and baryons. In contrast, for $N(1535)$ one needs some components other than hadronic composites, such as quark originated states. This twofold character of $N(1535)$, meson cloud and valence quark, can be seen also in the transition form factors of $\gamma^* N \rightarrow N(1535)$ (helicity amplitudes) [21]. In Ref. [21], the $N(1535)$ transition form factors obtained by two different approaches, the chiral unitary approach [22] and the spectator quark model [23,24], are compared. The quark model calculation of the transition form factors tells that the F_1^* form factor is produced well, while the F_2^* amplitude is overestimated in higher Q^2 , where the quark model may be applicable. It is very interesting that the meson cloud components calculated by the chiral unitary model compensates the disagreement of the quark model calculation as seen Fig. 1.

In the chiral unitary approach, one can calculate the diagonal component of the electromagnetic form factor, and in Ref. [25] the first moment of the $\Lambda(1405)$ form factor, which corresponds to the spatial radius if the particle is stable, and it has been found that $\Lambda(1405)$ has a substantially large size compared to the typical hadron. The “radius” of unstable particles are obtained as a complex number and its interpretation should be done carefully.

5 Kaonic few-body systems

Let us consider hadronic composite states with kaon and nucleon constituents as an extension of the $\Lambda(1405)$ resonance of a quasibound state of $\bar{K}N$. For hadronic composite states, pion has a too light mass to form bound states with other hadrons by hadronic interaction, since the pion kinetic energy in a hadronic composite system overcomes attractive potential energy. In contrast, kaon has a unique feature in hadronic composite state. The mass of kaon is so moderately heavy that kaon kinetic energy in hadronic bound systems can be smaller. In kaonic few-body systems, hadronic molecular states are unavoidably resonances decaying into pionic channels. The interaction of kaon as a Nambu-




Fig. 2 Family of kaonic few-body states.

Goldstone boson is described well chiral effective theory. It suggests that s -wave interactions in the $\bar{K}N$ and $\bar{K}K$ channels are attractive enough to form two-body quasibound states. The Tomozawa-Weinberg interaction is a driving force of the hadronic molecular system. It is very interesting that the strength of the Tomozawa-Weinberg interaction is fixed by SU(3) flavor symmetry and K and N are classified into the same state vector in the SU(3) octet representation. Therefore, considering also that K and N are a similar mass, one finds that the fundamental interactions in s -wave are very similar in the $\bar{K}K$ and $\bar{K}N$ channel. Consequently these channels with $I = 0$ have quasibound states of $\bar{K}K$ and $\bar{K}N$ with a dozen MeV binding energy. This similarity between K and N is responsible for systematics of three-body kaonic systems, $\bar{K}NN$, $\bar{K}KN$, $\bar{K}\bar{K}N$ and $\bar{K}KK$, as shown Fig. 2. Further few-body systems with kaons and nucleons have been studied, for instance, in $\bar{K}NNN$ and $\bar{K}\bar{K}NN$ systems [26, 27].

The $\bar{K}KN$ and $\bar{K}\bar{K}N$ states with $I = 1/2$ and $J^P = 1/2^+$, which are N^* and Ξ^* resonances, respectively, were studied first in Refs. [28, 29] with a single-channel non-relativistic potential model. The $\bar{K}KN$ system was found to be bound with 20 MeV binding energy [28], and later was investigated in a more sophisticated calculation [30, 31] based on a coupled-channels Faddeev method developed in Refs. [32], in which a very similar state to one obtained in the potential model was found. It was found also in a fixed center approximation of three-body Faddeev calculation [33]. The $\bar{K}KN$ state is essentially described by a coexistence of $K\Lambda(1405)$ and $f_0(980)N$ [28]. An experimental search for $\bar{K}KN$ was discussed in Ref. [34]. The $\bar{K}KK$ state with $I = 1/2$ and $J^P = 0^-$, being an excited state of kaon, was studied in a two-body f_0K and a_0K dynamics [35], in the three-body Faddeev calculation [36] and in the non-relativistic potential model [36]. The three-body Faddeev calculation was done in coupled-channels of $\bar{K}KK$, $K\pi\pi$ and $K\pi\eta$ and a resonance state was found at 1420 MeV, while the potential model suggested a quasi bound state with a binding energy 20 MeV. This state is essentially described by the $\bar{K}KK$ single channel and its configuration is found to be mostly f_0K . Experimentally, Particle Data Group tells that there is a excited kaon around 1460 MeV observed in $K\pi\pi$ partial-wave analysis, although it is omitted from the summary table.

In the potential model calculations of the $\bar{K}KN$ and $\bar{K}KK$ states, it was found that the root mean-squared radii of these systems are as large as 1.7 fm, which are similar with the radius of ^4He . The inter-hadron distances are comparable with an average nucleon-nucleon distance in nuclei. It was also found that the two-body subsystems inside the three-body bound state keep their properties in isolated two-body systems. These features are caused by weakly binding of the three hadrons.

6 Conclusion

There should be hadronic composite states in which hadrons including mesons are constituents of the state in hadron spectrum as one of the existence forms of hadrons. These states have spatially larger sizes than the typical confinement range, because the driving force of the hadronic composite state is inter-hadron interaction which is out of the confinement range.

The $\Lambda(1405)$ resonance is one of the strong candidates of the hadronic composite states. The peculiarity of the $\Lambda(1405)$ resonance is not only being a hadronic composite object but also being composed of two resonance states. These two states stems from the presence of two attractive channels in fundamental meson-baryon interactions. Eventually the observed $\Lambda(1405)$ resonance is composed by two pole states. One of the states is a quasi-bound state of $\bar{K}N$ located at around 1420 MeV and dominantly couples to the $\bar{K}N$ channel. Thus, this is the relevant resonance for the kaon-nucleus interaction. The double pole structure of $\Lambda(1405)$ can be confirmed experimentally by observing $K^-d \rightarrow \Lambda(1405)n$, in which $\Lambda(1405)$ is produced selectively by the $\bar{K}N$ channel and the peak position appears around 1420 MeV.

The idea that $\Lambda(1405)$ is a quasibound state of $\bar{K}N$ can be extended systematically to further few-body states with kaons like $\bar{K}NN$, $\bar{K}KN$ and $\bar{K}KK$ having dozens MeV binding energy. In these states, a unique role of kaon is responsible for the systematics of the few-body kaonic states. Kaon has a half mass of nucleon and a very similar coupling nature to nucleon in the s -wave chiral interaction. This leads to weakly bound systems within the hadronic interaction range. The hadronic composite state is a concept of weakly binding systems of hadron constituents. If a resonance state has a large binding energy measured from the break-up threshold, coupled-channel effects, like $\pi\Sigma$ against $\bar{K}N$, and/or shorter range quark dynamics should be important for the resonance state. In such a case the hadronic composite picture is broken down, and one should take into account coupled channels contributions and quark dynamics.

The hadronic composite configuration is a complementary picture of hadron structure to constituent quarks, which successfully describe the structure of the low-lying baryons in a simple way. Strong diquark configurations inside hadrons can be effective constituents [37], and mixture of hadronic molecular and quark originated states is also probable in some hadronic resonances [38]. The hadronic molecular state has a larger spacial size than the typical low-lying hadrons. In heavy ion collision, coalescence of hadrons to produce loosely bound hadronic molecular systems is more probable than quark coalescence for compact multi-quark systems [39,40]. Thus, one could extract the structure of hadrons by observing the production rate in heavy ion collisions.

Acknowledgements The author would like to acknowledge his collaborators of this work, T. Hyodo, A. Hosaka, A. Martínez Toress, Y. Kanada-En'yo, E. Oset, T. Sekihara, J. Yamagata-Sekihara, J.A. Oller, A. Ramos, U.-G. Meissner, M. Doring, G. Ramalho, K. Tsushima. This work was partially supported by the Grants-in-Aid for Scientific Research (No. 24105706), and done in part under the Yukawa International Program for Quark-hadron Sciences (YIPQS).

References

1. Dalitz, R.H., Tuan, S.F.: A possible resonant state in pion-hyperon scattering. *Phys. Rev. Lett.* **2**, 425–428 (1959). DOI 10.1103/PhysRevLett.2.425
2. Dalitz, R.H., Tuan, S.F.: The phenomenological representation of \bar{K} -nucleon scattering and reaction amplitudes. *Ann. Phys.* **10**, 307–351 (1960). DOI 10.1016/0003-4916(60)90001-4
3. Dalitz, R.H., Wong, T.C., Rajasekaran, G.: Model calculation for $Y_0^*(1405)$ resonance state. *Phys. Rev.* **153**, 1617–1623 (1967)
4. Veit, E.A., Jennings, B.K., Thomas, A.W., Barrett, R.C.: S -wave meson-nucleon scattering in an SU(3) cloudy bag model. *Phys. Rev.* **D31**, 1033 (1985). DOI 10.1103/PhysRevD.31.1033

5. Kaiser, N., Siegel, P.B., Weise, W.: Chiral dynamics and the low-energy kaon - nucleon interaction. Nucl. Phys. **A594**, 325–345 (1995). DOI 10.1016/0375-9474(95)00362-5
6. Hyodo, T., Jido, D.: The nature of the $\Lambda(1405)$ resonance in chiral dynamics. Prog. Part. Nucl. Phys. **67**, 55–98 (2012). DOI 10.1016/j.ppnp.2011.07.002
7. Ikeda, Y., Hyodo, T., Weise, W.: Improved constraints on chiral SU(3) dynamics from kaonic hydrogen. Phys. Lett. **B706**, 63–67 (2011). DOI 10.1016/j.physletb.2011.10.068
8. Ikeda, Y., Hyodo, T., Weise, W.: Chiral SU(3) theory of antikaon-nucleon interactions with improved threshold constraints. Nucl. Phys. **A881**, 98–114 (2012). DOI 10.1016/j.nuclphysa.2012.01.029
9. Bazzi, M., et al.: A New Measurement of Kaonic Hydrogen X rays. Phys. Lett. **B704**, 113–117 (2011). DOI 10.1016/j.physletb.2011.09.011
10. Oller, J.A., Meissner, U.G.: Chiral dynamics in the presence of bound states: kaon nucleon interactions revisited. Phys. Lett. **B500**, 263–272 (2001)
11. Jido, D., Oller, J.A., Oset, E., Ramos, A., Meissner, U.G.: Chiral dynamics of the two $\Lambda(1405)$ states. Nucl. Phys. **A725**, 181–200 (2003). DOI 10.1016/S0375-9474(03)01598-7
12. Hyodo, T., Weise, W.: Effective $\bar{K}N$ interaction based on chiral SU(3) dynamics. Phys. Rev. **C77**, 035204 (2008). DOI 10.1103/PhysRevC.77.035204
13. Jido, D., Oset, E., Sekihara, T.: Kaonic production of $\Lambda(1405)$ off deuteron target in chiral dynamics. Eur. Phys. J. **A42**, 257–268 (2009). DOI 10.1140/epja/i2009-10875-5
14. Jido, D., Oset, E., Sekihara, T.: Kaon induced $\Lambda(1405)$ production on a deuteron target at DAFNE. Eur. Phys. J. **A47**, 42 (2011). DOI 10.1140/epja/i2011-11042-3
15. Noumi, H., et al.: J-PARC proposal E31 “Spectroscopic study of hyperon resonances below $\bar{K}N$ threshold via the (K^-, n) reaction on Deuteron” (2009)
16. Weinberg, S.: Evidence that the deuteron is not an elementary particle. Phys. Rev. **137**, B672–B678 (1965)
17. Lurié, D., Macfarlane, A.J.: Equivalence Between Four-Fermion and Yukawa Coupling, and the $Z_3 = 0$ Condition for Composite Bosons. Phys. Rev. **136**, B816–B829 (1964). DOI 10.1103/PhysRev.136.B816
18. Hyodo, T., Jido, D., Hosaka, A.: Compositeness of dynamically generated states in a chiral unitary approach. Phys. Rev. C **85**, 015201 (2012). DOI 10.1103/PhysRevC.85.015201
19. Hyodo, T., Jido, D., Hosaka, A.: Origin of the resonances in the chiral unitary approach. Phys. Rev. **C78**, 025203 (2008). DOI 10.1103/PhysRevC.78.025203
20. Ecker, G., Gasser, J., Pich, A., de Rafael, E.: The role of resonances in chiral perturbation theory. Nucl. Phys. **B321**, 311–342 (1989)
21. Ramalho, G., Jido, D., Tsushima, K.: Valence quark and meson cloud contributions for the $\gamma^* \Lambda \rightarrow \Lambda^*$ and $\gamma^* \Sigma_0 \rightarrow \Lambda^*$ reactions. Phys. Rev. D **85**, 093014 (2012). DOI 10.1103/PhysRevD.85.093014
22. Jido, D., Doering, M., Oset, E.: Transition form factors of the $N^*(1535)$ as a dynamically generated resonance. Phys. Rev. **C77**, 065207 (2008). DOI 10.1103/PhysRevC.77.065207
23. Ramalho, G., Pena, M.T.: A covariant model for the $\gamma N \rightarrow N(1535)$ transition at high momentum transfer. Phys. Rev. **D84**, 033007 (2011). DOI 10.1103/PhysRevD.84.033007. [Publisher-note D84:039904,2011]
24. Ramalho, G., Tsushima, K.: A simple relation between the $\gamma N \rightarrow N(1535)$ helicity amplitudes. Phys. Rev. **D84**, 051301 (2011). DOI 10.1103/PhysRevD.84.051301
25. Sekihara, T., Hyodo, T., Jido, D.: Internal structure of the resonant $\Lambda(1405)$ state in chiral dynamics. Phys. Rev. C **83**(5), 055202 (2011). DOI 10.1103/PhysRevC.83.055202
26. Akaishi, Y., Yamazaki, T.: Nuclear \bar{K} bound states in light nuclei. Phys. Rev. **C65**, 044005 (2002). DOI 10.1103/PhysRevC.65.044005
27. Barnea, N., Gal, A., Liverts, E.: Realistic calculations of $\bar{K}NN$, $\bar{K}N\bar{N}N$, and $\bar{K}\bar{K}NN$ quasibound states. Phys. Lett. **B712**, 132–137 (2012). DOI 10.1016/j.physletb.2012.04.055
28. Jido, D., Kanada-En'yo, Y.: $K\bar{K}N$ molecule state with $I = 1/2$ and $J^P = 1/2^+$ studied with three-body calculation. Phys. Rev. **C78**, 035203 (2008). DOI 10.1103/PhysRevC.78.035203
29. Kanada-En'yo, Y., Jido, D.: $\bar{K}\bar{K}N$ molecule state in three-body calculation. Phys. Rev. C **78**, 025212 (2008). DOI 10.1103/PhysRevC.78.025212
30. Martinez Torres, A., Khemchandani, K.P., Oset, E.: Solution to Faddeev equations with two-body experimental amplitudes as input and application to $J^P = 1/2^+$, $S = 0$ baryon resonances. Phys. Rev. **C79**, 065207 (2009). DOI 10.1103/PhysRevC.79.065207
31. Martinez Torres, A., Jido, D.: $K\Lambda(1405)$ configuration of the $K\bar{K}N$ system. Phys. Rev. **C82**, 038202 (2010). DOI 10.1103/PhysRevC.82.038202

32. Martinez Torres, A., Khemchandani, K.P., Oset, E.: Three body resonances in two meson-one baryon systems. *Phys. Rev. C* **77**, 042203 (2008). DOI 10.1103/PhysRevC.77.042203
33. Xie, J.J., Martinez Torres, A., Oset, E.: Faddeev fixed center approximation to the $N\bar{K}K$ system and the signature of a $N^*(1920) (1/2^+)$ state. *Phys. Rev. C* **83**, 065207 (2011). DOI 10.1103/PhysRevC.83.065207
34. Martinez Torres, A., Khemchandani, K.P., Meissner, U.G., Oset, E.: Searching for signatures around 1920 MeV of a N^* state of three hadron nature. *Eur. Phys. J. A* **41**, 361–368 (2009). DOI 10.1140/epja/i2009-10834-2
35. Albaladejo, M., Oller, J.A., Roca, L.: Dynamical generation of pseudoscalar resonances. *Phys. Rev. D* **82**, 094,019 (2010). DOI 10.1103/PhysRevD.82.094019
36. Martinez Torres, A., Jido, D., Kanada-En'yo, Y.: Theoretical study of the $KK\bar{K}$ system and dynamical generation of the $K(1460)$ resonance. *Phys. Rev. C* **83**, 065205 (2011). DOI 10.1103/PhysRevC.83.065205
37. Kim, K., Jido, D., Lee, S.H.: Diquarks: a QCD sum rule perspective. *Phys. Rev. C* **84**, 025204 (2011). DOI 10.1103/PhysRevC.84.025204
38. Nagahiro, H., Nawa, K., Ozaki, S., Jido, D., Hosaka, A.: Composite and elementary natures of $a_1(1260)$ meson. *Phys. Rev. D* **83**(11), 111504 (2011). DOI 10.1103/PhysRevD.83.111504
39. Cho, S., et al.: Identifying multiquark hadrons from heavy ion collisions. *Phys. Rev. Lett.* **106**(21), 212001 (2011). DOI 10.1103/PhysRevLett.106.212001
40. Cho, S., et al.: Exotic Hadrons in Heavy Ion Collisions. *Phys. Rev. C* **84**, 064910 (2011). DOI 10.1103/PhysRevC.84.064910